

ISOMETRY TYPES OF PROFINITE GROUPS

A.IVANOV (WROCLAW)

Abstract. ^{1 2 3} Let T be a rooted tree and $Iso(T)$ be the group of its isometries. We study closed subgroups G of $Iso(T)$ with respect to the number of conjugacy classes of $Iso(T)$ having representatives in G .

1. INTRODUCTION

Let T be an infinite locally finite tree with the root $\emptyset$. By $layer_n$ we denote the set of all elements of T , n -distant from $\emptyset$ (i.e. elements of level n). We always assume that there is an infinite sequence $\bar{m} = m_1, m_2, \dots$ of natural numbers such that T is of the form $T_{\bar{m}}$, the tree where all elements of $layer_n$ are of valency $m_{n+1} + 1$ for $n > 0$. Let $T_n = \bigcup_{i \leq n} layer_i$ and $T_{\langle n \rangle}$ be the tree of the isomorphism type of T_w , the tree hanging from $w \in layer_n$.

The group $Iso(T)$ of all isometries of T fixing $\emptyset$ is a profinite group with respect to canonical homomorphisms $\pi_n : Iso(T) \rightarrow Iso(T_n)$. In fact the following examples represent all the main fields in the subject.

Consider the group $GL_d(\mathbb{Z}_p)$ and its action on $D = (\mathbb{Z}_p)^d$. Let T_{ad} be the tree of all cosets in $(\mathbb{Z}_p)^d$ with respect to all subgroups of the form $p^k(\mathbb{Z}_p)^d$. If G is a closed subgroup of $GL_d(\mathbb{Z}_p)$, then D becomes a continuous $G\mathbb{Z}_p$ -module. Thus G acts by isometries on T_{ad} and fixes the root D . A very close example is the action of $PGL_2(\mathbb{Z}_p)$ on the tree of lattices in $\mathbb{Q}_p \times \mathbb{Q}_p$ (see [18]).

Completely different groups are provided by Grigorchuk-type constructions. In [13] *branch groups* are introduced, which in some sense can be considered as an

¹**2000 Mathematics Subject Classification:** 20A15, 03C45, 20E08, 20E18, 20E45

²**Key words and phrases:** rooted trees, isometry groups, profinite groups, conjugacy classes, interpretability

³the research was supported by KBN grant 1 P03A 025 28

axiomatization of well-known examples (their profinite completions) of Burnside groups found by Grigorchuk.

Are there any global (=model-theoretic) properties giving interesting dividing lines in these examples ?

In this paper we concentrate on characterizations concerning diversity of elements occurring in the group. Our basic property can be described as follows. Consider the relation $\sim$ of conjugacy in the group $Iso(T)$ of all isometries of T . It is easy to see that every conjugacy class is a closed subset of $Iso(T)$. Let G be a closed subgroup of $Iso(T)$. We say that G has a *small (resp. countable) number of isometry types* if G meets only $< 2^{\aleph_0}$ (countably many) conjugacy classes of $Iso(T)$. Since $G/\sim$ is analytic, smallness is equivalent to countability.

We study profinite groups which can be embedded into $Iso(T)$ as closed subgroups with a small number of isometry types. It seems to us that this class naturally arises in many situations. One of them has model-theoretic flavor and is worth mentioning here. For closed $G < Iso(T)$ consider the corresponding inverse system $G_0 \leftarrow G_1 \leftarrow \dots \leftarrow G_i \leftarrow \dots$ with $G_i \leq Iso(T_i)$. Any automorphism of G fixing all G_i (as sets) is called a *profinite automorphism* of G . By $Aut^*(G)$ we denote the group of all profinite automorphisms.

Assume that any profinite automorphism of G is induced by an isometry of T as an inner automorphism of $Iso(T)$ (by Proposition 8.1 from [15] this happens if for any $w \in T$ the point-wise stabilizer of $T \setminus T_w$ is level-transitive on T_w). If the *number of $Aut^*(G)$ -orbits on G is at most countable* then G has a small number of isometry types. The former condition has been introduced by Newelski in [16] as a counterpart of small theories from model theory. The main (still open) conjecture in these respects states that profinite groups with a small number of $Aut^*(G)$ -orbits have open abelian subgroups. Some partial confirmation has been obtained by F.Wagner in [21].

In fact Newelski's conjecture has become the first motivation for our interests to isometry groups with a small number of isometry types. Notice that when G is an inverse limit of a system of finite groups $G_0 \leftarrow G_1 \leftarrow \dots \leftarrow G_i \leftarrow \dots$ with kernels N_i of the corresponding canonical homomorphisms $G \rightarrow G_i$, then G acts by left multiplication on the tree $T_C(G)$ of left cosets of all N_i , $i \in \omega$ (any edge of

the tree is defined by a pair of the form $gN_{i+1} \subset gN_i$. In this case any profinite automorphism of G (which stabilises all N_i) becomes an isometry of $T_C(G)$. Thus if G has a small number of $\text{Aut}^*(G)$ -orbits, then G has a small number of isometry types with respect to this tree action.

It has become very surprising for us that the smallness condition above has very strong abstract consequences concerning the structure of G . In particular it turns out that we can say quite a lot about G when G is nilpotent.

We also study the related property of definability of the action in $\mathbb{Q}_p$ and show that some natural zeta functions arising in these respects are rational. We compare our approach with the one suggested by Z.Charzidakis in [5]. We will see that ω -categoricity appearing in [5] is very far from smallness of the number of isometry types (at least in the case of examples above).

1.1. Preliminaries. Let $T = T_{\bar{m}}$, where $\bar{m}$ is an infinite sequence of positive natural numbers. For every n we define an alphabet $A_n = \{a_1^n, \dots, a_{m_n}^n\}$ and consider T as the union of all finite products $A_1 \times \dots \times A_n$, $n \in \omega$, interpreted as paths of T started at $\emptyset$: here we identify layer_n with $A_1 \times \dots \times A_n$

If v is a vertex of T and $g \in \text{Iso}(T)$, then by $g(v)$ we usually denote the result of g applied to v . On the other hand we sometimes use the form v^g , because it becomes more convenient when one considers iterated wreath products. Then as in [13] any $g \in \text{Iso}(T)$ is regarded as a labelling $\{\gamma(v)\}_{v \in T}$ of the vertices by elements of the symmetric group which acts on the edges below the vertex. The effect of $g \in \text{Iso}(T)$ on a vertex u corresponding to $p_u = (a_1, \dots, a_n)$, a path from $\emptyset$ with $a_i \in A_i$, is given by the formula:

$$p_u^g = (a_1^{\gamma(\emptyset)}, a_2^{\gamma(a_1)}, \dots, a_n^{\gamma(a_1 \dots a_{n-1})}),$$

where by $a^{\gamma(v)}$ we denote the result of the permutation $\gamma(v)$ applied to the edge a . The rule for the composing automorphisms is given by the following formulas (see [13]):

Let $h = \{\delta(v)\}_{v \in T}$, $f = \{\sigma(v)\}_{v \in T}$ and $g = \{\gamma(v)\}_{v \in T}$.

If $h = fg$, then $\delta(v) = \sigma(v)\gamma(v^f)$.

If $f = g^{-1}$, then $\sigma(v) = (\gamma(v^{g^{-1}}))^{-1}$.

Now assume that for every n we have chosen a permutation group $P_i \leq \text{Sym}(A_i)$. Then the subgroup of all labellings $\{\gamma(v)\}_{v \in T}$ with $\gamma(v) \in P_{i+1}$ for $v \in \text{layer}_i$ is called *the iterated wreath product* of (P_i, A_i) , $i \in \{1, 2, \dots\}$ and is denoted by $\lambda_\omega(P_i, A_i)$.

With any $v \in T$ (an end $\mathbf{e} \in \partial T$ resp.) and a group $G \leq \text{Iso}(T)$ we associate the stabilizer $St_G(v)$ ($St_G(\mathbf{e})$ resp.). The level stabilizer is defined by $St_G(n) = \bigcap \{St_G(v) : |p_v| = n\}$. It is clear that $St_G(n) = \text{Ker} \pi_n$, where $\pi_n : \text{Iso}(T) \rightarrow \text{Iso}(T_n)$ is defined as above. On the other hand $St_G(n)$ is naturally identified with a subgroup of $\prod \{Iso(T_v) : v \in \text{layer}_n\}$.

The *right stabilizer* of v is defined to be $Rs_G(v) = \{g \in \text{Iso}(T) : g \text{ fixes } T \setminus T_v \text{ pointwise}\}$. The right stabilizer of the n -th level $Rs_G(n)$ is defined to be the group generated by $\bigcup \{Rs_G(v) : v \in \text{layer}_n\}$. It is clear that $Rs_G(n) = \oplus \{Rs_G(v) : v \in \text{layer}_n\}$.

We now define (weakly) branch groups. They have been introduced by R.Grigorchuk in [13]. They are one of the main objects in the area.

Let T be a locally finite tree, $T = T_{\bar{m}}$. A closed subgroup $G < \text{Iso}(T)$ is called a branch group with respect to T if

- (i) G acts transitively on each layer of the tree, and
- (ii) for each $n \geq 1$ there exists closed $L_n \leq \text{Iso}(T_{\langle n \rangle})$ such that the direct product $H_n = L_n^{\text{layer}_n}$ is normal and of finite index in G (each factor of the product acts on the subtree hanging from the corresponding vertex of the n -th layer).

Condition (ii) of this definition is equivalent to $|G : Rs_G(n)| < \infty$ for every $n \in \omega$ [13]. If we replace (ii) by the condition that $Rs(n)$ is always infinite, then we obtain a definition of *weakly branch groups*.

Let $g \in \text{Iso}(T)$. By T_g we denote the *orbit tree* of g , i.e. the set of all g -cycles $T/\langle g \rangle$ with respect to the natural adjacency. The orbit tree is labelled by natural numbers as follows: to each orbit $\langle g \rangle v$ we associate the size $|\langle g \rangle v|$.

The following theorem from [10] will be one of the main tools of the paper:

isometries g and h are conjugated in $\text{Iso}(T)$ if and only if the corresponding orbit trees T_g and T_h are isomorphic by a labelling preserving isomorphism.

It is worth noting that the class of groups having a faithful representation with a small number of isometry types is closed under direct sums: if G_1 and G_2 have appropriate realisations on trees of valencies N_1 and N_2 , then $G_1 \oplus G_2$ can be realised on a tree of valency $N_1 + N_2$ so that the number of isometry types is small.

1.2. Main results. In Section 2 we study profinite groups which can be realized as closed subgroups of $Iso(T)$ which have a small number of isometry types. We show that $\mathbb{Z}_p$ and the direct power H^ω of a finite group H have such representations in $Iso(T)$ for appropriate T . We conjecture that nilpotent profinite groups G with this property are very close to these examples. In Section 2.1 we slightly confirm this showing that only finitely many primes can divide $|G|$ and the periodic part of G is of finite exponent.

We show that the smallness condition is so restrictive itself that for example weakly branch groups do not have a small number of isometry types. We will also see here that the tree of cosets of kernels of canonical projections naturally arises in these respects, and the smallness of the number of isometry types implies some interesting properties of this tree (Proposition 2.2).

In Section 2.2 we study pronilpotent groups G which have a small number of isometry types for all faithful continuous representations in isometry groups of locally finite trees. In the case when the group G is nilpotent we show that such a group must be of finite rank. This condition becomes sufficient when the group is abelian (see Theorem 2.9).

In Section 3 we study another condition of smallness. Let T be a locally finite rooted tree. For every natural n we introduce the following equivalence relation on $Iso(T)$: for $g, g' \in Iso(T)$ we define $g \equiv_n g'$ if the restrictions of g and g' to T_n are conjugated in $Iso(T_n)$.

For a subgroup $G \leq Iso(T)$ we denote by c_n^G the number of all $\equiv_n$ -classes meeting G . In Section 3 we will study the question when the zeta function $\sum_{n \geq 0} c_n^G t^n$ is rational.

In general one cannot expect, that this happens for a typical $G < Iso(T)$ For example Grigorchuk shows in [12] that the Hilbert-Poincaré series of some of his

groups are not rational. But if we smooth up the situation (under some model theoretic assumptions) we can have very reasonable variants of the question. Moreover examples of the introduction with $GL_d(\mathbb{Z}_p)$ involved become under our methods.

The idea looks as follows. Let Γ be a closed subgroup of $Iso(T)$. We say that $g, g' \in Iso(T)$ have the same Γ -type if there exists $h \in \Gamma$ such that $g^h = g'$. It makes sense to study Γ -types of $Iso(T)$ assuming that Γ is interpretable in some natural model-theoretic objects (not only in the case $\Gamma = Iso(T)$). In this situation we say that $G \leq Iso(T)$ has a small number of Γ -types if G meets $\leq \aleph_0$ Γ -orbits of the conjugacy action of Γ on $Iso(T)$.

For every natural n we introduce the following equivalence relation on $Iso(T)$: for $g, g' \in Iso(T)$ we define $g \equiv_n^\Gamma g'$ if the restrictions of g and g' to T_n are conjugated by an element of the projection of Γ to T_n . Now for the subgroup $G \leq Iso(T)$ we denote by c_n^G the number of all $\equiv_n^\Gamma$ -classes meeting G . We will really study the question when the zeta function $\sum_{n \geq 0} c_n^G t^n$ is rational for some Γ interpretable in the field $\mathbb{Q}_p$. In Section 3 we discuss a number of examples. In particular we study a point-stabilizer of the action of $SL_2(\mathbb{Q}_p)$ on the tree of lattices [18]. In fact our material here is based on a nice paper of E.Hrushovski and B.Martin [9].

We must mention that in the eighties Z.Chatzidakis introduced some language for profinite groups and showed that model-theoretic investigations of the corresponding structures (denoted by $S(G)$) give very strong consequences in purely algebraic questions about profinite groups and field extensions. Moreover structures $S(G)$ arising in this approach, have very nice model-theoretic properties. In some typical situations they are ω -categorical, i.e. small. It should be natural to start model-theoretic investigations of closed subgroups of $Iso(T)$ with these aspects. We fill this gap in Section 4. The answer which we obtain here, shows that the majority of our examples represent the simplest case of this classification. In fact we show that the dividing line of ω -categoricity of structures $S(G)$ is situated inside the class of branch groups. We give here some interesting examples. In particular the profinite completion of the famous example of Grigorchuk is ω -categorical in this sense. These results show why the condition of smallness which comes from the Newelski's approach is more appropriate for our examples.

2. GROUPS WITH A SMALL NUMBER OF ISOMETRY TYPES

2.1. Groups having representations with a small number of isometry types. We now consider groups which can be realized as closed subgroups of $Iso(T)$ with a small number of isometry types. Let T be a locally finite tree and G be a closed subgroup of $Iso(T)$. By π_n we denote the canonical homomorphism from G to the group G_n of all restrictions of G to T_n , $n \in \omega$.

Lemma 2.1. (1) *If there is some $n \in \omega$ such that $\ker(\pi_n) \setminus \{1\}$ consists of isometries of the same type, then the group G has an open subgroup which is a locally finite pro- p -group for some prime p .*

(2) *If there is $n \in \omega$ such that the set of non-diagonal pairs of $\ker(\pi_n)$ is contained in the same orbit of $Iso(T)$, then $\ker(\pi_n)$ is isomorphic to the power $(\mathbb{Z}(p))^\omega$.*

Proof. Under the assumptions of the lemma we easily see that for every $l \geq n$ all non-trivial elements of the finite group $\ker(\pi_n)/\ker(\pi_l)$ are of the same order. Thus all of them are of order p , where p is prime. This shows that $\ker(\pi_n)$ is a pro- p -group of exponent p . By a theorem of Zelmanov from [22] it is locally finite.

The second statement follows from the fact that every $\ker(\pi_n)/\ker(\pi_l)$ is a nilpotent group and has non-trivial centre. The assumption says that it must be isomorphic to a finite power of $\mathbb{Z}(p)$. The rest is obvious. $\square$

Proposition 2.2. *Let T be a locally finite tree and G be a closed subgroup of $Iso(T)$ with a small number of isometry types. Then the following statements hold.*

(1) *For every $m \in \omega$ and every $h \in G$ there are some $n \in \omega$ and $g \in G \setminus \ker(\pi_n)$ such that $g \cdot \ker(\pi_n) \subseteq h \cdot \ker(\pi_m)$ and $g \cdot \ker(\pi_n)$ consists of isometries of the same type.*

(2) *If $g \in G \setminus \ker(\pi_n)$, $n \in \omega$, and the set of all non-diagonal pairs of $g \cdot \ker(\pi_n)$ is contained in the same orbit of $Iso(T)$, then G has an open subgroup which is a locally finite pro- p -group for some prime p . If moreover for every $g_1, g_2, g'_1, g'_2 \in g \cdot \ker(\pi_n)$ with $g_1 \neq g_2$, $g'_1 \neq g'_2$ one of the pair (g_1, g_2) and (g'_1, g'_2) can be mapped to another by an isometry fixing some element of $g \cdot \ker(\pi_n)$ then $\ker(\pi_n)$ is isomorphic to the power $(\mathbb{Z}(p))^\omega$, where p is prime.*

Proof. (1) By the Baire category theorem, $Iso(T)$ -conjugacy classes which are open in G form a dense subset of G . By the definition of the profinite topology we see the existence of $g \cdot \ker(\pi_n)$ as in the formulation.

(2) If the set of all non-diagonal pairs of $g \cdot \ker(\pi_n)$ is contained in the same orbit of $Iso(T)$, then each element of $\ker(\pi_n)$ can be mapped to any other element of $\ker(\pi_n)$ by an isometry. By Lemma 2.1, G has an open subgroup which is a locally finite pro- p -group for some prime p .

Assuming the last condition of the formulation consider any $\ker(\pi_n)/\ker(\pi_l)$ with $l \geq n$. As we already know this is a group of exponent p . Let $h_1 \cdot \ker(\pi_l)$ belong to the centre of $\ker(\pi_n)/\ker(\pi_l)$, and $h_2 \neq h_3 \in \ker(\pi_n)$. Let an isometry μ map gh_1 to gh_2 and fix the pair g, gh_3 (we may assume so). Then we have modulo $\ker(\pi_l)$: $gh_1 \cdot gh_3 = ggh_3(h_1)^g = g(h_3)^{g^{-1}} \cdot g(h_1)^g$ and $gg(h_2)^g h_3 = gh_2gh_3 = \mu(gh_1)\mu(gh_3) = \mu(gh_1 \cdot gh_3) = g(h_3)^{g^{-1}}g(h_2)^g = ggh_3(h_2)^g$. Thus $h_2\ker(\pi_l)$ belongs to the centre of $\ker(\pi_n)/\ker(\pi_l)$. The rest is obvious. $\square$

Corollary 2.3. *Let T be a locally finite tree and G be a closed subgroup of $Iso(T)$ with a small number of isometry types. Then there is no infinite sequence of vertices $v_1, v_2, \dots$ such that $T_{v_i} \cap T_{v_j} = \emptyset$ for $i \neq j$, and for every i the point-wise stabilizer of $T \setminus T_{v_i}$ in G has a non-trivial action on T_{v_i} . In particular a weakly branch group does not have a small number of isometry types.*

Proof. Assume the contrary. Since T is locally finite, we easily see that G does not satisfy Proposition 2.2 (1). The rest is obvious. $\square$

Example. One can build a tree action of the wreath product $\mathbb{Z}(p)wr\mathbb{Z}_p$ with a large number of isometry types.

The following example concerns the case of a tree of valency three.

Example. Let $T = \{0, 1, \dots\}^{<\mathbb{N}}$ be a tree with the root $\emptyset$ ($\mathbb{N} = \omega \setminus \{0\}$; the elements of the tree are presented by finite sequences). Let $g \in Iso(T)$ be defined as follows: $g(0) = 1$, $g(1) = 0$. We define g so that at every level g has a unique q -cycle: If v consists of n 0-s (the beginning of the cycle) then let $g^i(v0) = g^i(v)0$, $i < 2^n$, and $g^i(v0) = g^i(v)1$, $2^n \leq i < 2^{n+1}$. It is clear that g realises the *adding*

machine: the result of the application of g to a branch $(a_1, a_2, \dots)$ is the 2-adic sum $(a_1, a_2, \dots) + (1, 0, \dots)$. Let $G = cl(\langle g \rangle)$; then $G \cong \mathbf{Z}_2$, the additive group of 2-adic integers.

If $g' \in G \cap (Ker\pi_n \setminus Ker\pi_{n+1})$ then for every level $m > n$ there exists exactly 2^n g' -cycles of length 2^{m-n} . This follows from the fact that g' can be interpreted as a 2-adic number $c_{n+1}2^{n+1} + c_{n+2}2^{n+2} + \dots$ with $c_{n+1} = 1$. (Thus $c_{n+1}2^{n+i+1} + c_{n+2}2^{n+i+2} + \dots$ corresponds to $(g')^{2^i}$.)

By the description of conjugacy in $Iso(T)$ given in [10] we see that g_1 and g_2 from G have the same isometry type if and only if there is a number $n \in \omega$ such that $g_1, g_2 \in Ker\pi_n \setminus Ker\pi_{n+1}$. This obviously implies that the group G has a small number of isometry types. $\square$

Theorem 2.4. *Let T be a locally finite rooted tree.*

(2) *If a pronilpotent group G has a faithful representation in $Iso(T)$ with a small number of isometry types, then the number of primes dividing $|G|$ is finite.*

(2) *If a nilpotent group G has a faithful representation in $Iso(T)$ with a small number of isometry types, then the periodic part of G is of finite exponent.*

Proof. (1) Let G be the inverse limit of the sequence $G_1 \leftarrow G_2 \leftarrow G_3 \leftarrow \dots$ corresponding to an appropriate representation on a tree. Let P be the set of primes dividing $|G|$. Since G is pronilpotent, for every $X \subseteq P$ there exists $g_X \in G$ such that a prime number p divides $|g_X|$ if and only if $p \in X$. If P is infinite there is continuum many elements of G having pairwise distinct orders in $Iso(T)$. Any two such elements cannot be conjugated in $Iso(T)$.

(2) Assume the contrary. Taking a subgroup if necessary we may assume that G is nilpotent and periodic. Note that if each factor of the upper central series is of finite exponent then G is of finite exponent. Thus we can find a characteristic subgroup $G^- < G$ of finite exponent such that the center $C(G/G^-)$ is infinite and is not of bounded exponent.

Let π_n be the standard projection $G \rightarrow G_n$ corresponding to the restriction to T_n . We now build a sequence of natural numbers $n_1 < n_2 < \dots < n_i, \dots$ and a sequence $g_1, \dots, g_i, \dots \in G \setminus G^-$ with $g_i G^- \in C(G/G^-)$ such that for all $i \in \omega \setminus \{0\}$ the following properties hold:

- (a) $\prod_{j \leq i} |g_j| < |g_{i+1}|$;
- (b) $|\pi_{n_i}(g_i)| = |g_i|$;
- (c) $g_{i+1} \in \text{Ker}(\pi_{n_i})$.

This can be done by induction as follows. Using the fact that $C(G/G^-)$ is infinite and is not of finite exponent we choose g_{i+1} such that conditions (a) and (c) are satisfied. Since $|g_{i+1}| < \infty$, we can find n_{i+1} such that (b) is satisfied.

Now for every subset $Y \subset \omega$ take $g_Y := \prod \{g_i : i \in Y\}$. It is clear that g_Y is the limit of all products $\prod \{g_j : j \leq i, j \in Y\}$. If $Y \neq Y'$ and i is the least natural number from $Y \Delta Y'$, then the restrictions of g_Y and $g_{Y'}$ to T_{n_i} have distinct orders and cannot be conjugated by an isometry. This contradicts the assumption that G has a small number of isometry types. $\square$

Note that it is not very difficult to realize the situation of Theorem 2.4(2). The following observation gives a wide class of examples of representations with small numbers of isometry types.

Proposition 2.5. *Let H be a finite group. Then the product $G = H^{\aleph_0}$ has a faithful action on a rooted tree with countably many isometry types.*

Proof. Consider the inverse system of projections

$$H \leftarrow H \oplus H \leftarrow H \oplus H \oplus H \leftarrow \dots$$

where the groups consist of sequences

$$H = \{h_1, h_2, \dots, h_n\}, H \oplus H = \{h_1 h_1, h_1 h_2, \dots, h_{n-1} h_n, h_n h_n\}$$

$$, \dots, H \oplus \dots \oplus H = \{h_1 \dots h_1, \dots, h_n \dots h_n\}, \dots$$

The same sequences form an n -ary tree where the groups act by multiplication: $h_i w \cdot h_l = h_j w$ for $h_i \cdot h_l = h_j$, $h_i h_m w \cdot h_k h_l = h_j h_n w$ for $(h_i, h_m) \cdot (h_k, h_l) = (h_j, h_n), \dots$. It is easy to see that if $g \in G$ and $|g| = s$ then there are sequences $n_1, n_2, \dots, n_t$ and $s_1, \dots, s_t = s$ such that $|\pi_{n_1-1}(g)| = 1$, $|\pi_{n_1}(g)| = s_1 = |\pi_{n_2-1}(g)|$, $|\pi_{n_2}(g)| = s_2 = |\pi_{n_3-1}(g)|, \dots$ and $|\pi_{n_t}(g)| = s_t = |\pi_m(g)|$ for all $m \geq n_t$. In that case for every level $n_j \leq m < n_{j+1}$ all g -cycles are of length s_j and for every level $m \geq n_t$ all g -cycles are of length s . By the description of conjugacy in $\text{Iso}(T)$ given in [GNS] we see that g_1 and g_2 from G have the same isometry type if and only

if the corresponding sequences for them are the same. This obviously implies that the group G has a countable number of isometry types. $\square$

We now see that any group of the form

$$\mathbb{Z}(n_1)^\omega \oplus \dots \oplus \mathbb{Z}(n_k)^\omega \oplus \mathbb{Z}_{p_1}^{k_1} \oplus \dots \oplus \mathbb{Z}_{p_l}^{k_l}$$

has a faithful representation with a small number of isometry types.

Conjecture. An abelian profinite group A has a faithful representation with a small number of isometry types if and only if

$$A \cong \mathbb{Z}(n_1)^\omega \oplus \dots \oplus \mathbb{Z}(n_k)^\omega \oplus \mathbb{Z}_{p_1}^{k_1} \oplus \dots \oplus \mathbb{Z}_{p_l}^{k_l}.$$

Proposition 2.2 motivates the question of description of subgroups G of $Iso(T)$ having a small number of isometry types in $G \times G$. We have a small remark concerning this case.

Lemma 2.6. *If $G < Iso(T)$ has a small number of isometry types of $G \times G$, then G does not have an element of infinite order.*

Proof. Indeed, if $|g| = \infty$, then for any $z_1, z_2 \in \mathbb{Z}_p$ with $z_1 \neq z_2$ we have $g^{z_1} \neq g^{z_2}$. If an isometry $h \in Iso(T)$ centralizes g , then it must centralize any g^z , $z \in \mathbb{Z}_p$. Thus all pairs (g, g^z) , $z \in \mathbb{Z}_p$, have pairwise distinct isometry types. $\square$

Remark. The group $A = \prod_\omega \langle g_i \rangle$ with $\langle g_i \rangle \cong \mathbb{Z}(p)$, is an example of a small group in the sense of Newelski (this was observed by G.Bezulski in his master thesis). It is worth noting that the action of A on a binary tree as in Proposition 2.5, has a small number of isometry types. We thus consider the inverse system of projections

$$\mathbb{Z}(2) \leftarrow \mathbb{Z}(2) \oplus \mathbb{Z}(2) \leftarrow \mathbb{Z}(2) \oplus \mathbb{Z}(2) \oplus \mathbb{Z}(2) \leftarrow \dots$$

where the groups consist of sequences $0, 1, 00, 01, 10, 11, 000, \dots, 111, \dots$. The same sequences form a binary tree where the groups act as follows: $0w + 0 = 0w$, $0w + 1 = 1w$, $00w + 00 = 00w$, $\dots$, $01w + 11 = 10w$, $11w + 11 = 00w, \dots$. It is easy to see that if $g' \in A \cap (Ker\pi_n \setminus Ker\pi_{n+1})$ then for every level $m > n$ there exists exactly 2^{m-n-1} g' -cycles of length 2. By the description of conjugacy in $Iso(T)$ given in [10] we see that g_1 and g_2 from A have the same isometry type if and only if there

is a number $n \in \omega$ such that $g_1, g_2 \in \text{Ker}\pi_n \setminus \text{Ker}\pi_{n+1}$. This obviously implies that the group A has a small number of isometry types.

On the other hand by Lemma 2.3 the group A does not have a small number of isometry types under the following presentation in $T = p^{\mathbb{N}}$: for any i the element g_i stabilizers $T \setminus T_{1\dots 10}$ and $(1\dots 10jw)g_i = 1\dots 10(j+1)w$. $\square$

2.2. Isometry groups having a small number of isometry types for all isometry representations.

Lemma 2.7. (1) *Let a profinite group G and a sequence $g_i \in G$, $i \in \omega$, satisfy the condition that for every $i \in \omega$ there is a continuous epimorphism ρ_i from G to a finite group such that $\rho_i(g_i) \neq 1$, but all g_j with $j \neq i$ belong to the kernel of ρ_i . Then there is a locally finite rooted tree T such that G has a faithful representation in $\text{Iso}(T)$ with continuum many isometry types.*

(2) *If G is a pro- p -group as above, then there is a faithful representation of G in $\text{Iso}(T)$ with continuum many isometry types, where T is the rooted tree of valency $p+1$.*

Proof. (1) We define T as a composition of two parts. To define the first one assume that G can be presented as an inverse limit of a system $G_0 \leftarrow G_1 \leftarrow \dots \leftarrow G_i \leftarrow \dots$ of finite groups. Let N_i , $i \in \omega$, be the kernels of the corresponding canonical homomorphisms $G \rightarrow G_i$. Then G acts by left multiplication on the tree $T_C(G)$ of left cosets of all N_i , $i \in \omega$.

The second part $T_S(G)$ of the tree consists of all elements of $\bigcup\{(\bigoplus_{i=0}^j \rho_i(G)) : j \in \omega\}$, where the ordering of the tree is defined by extension. We define an action of G on this tree as follows. For every $g \in G$ and every vertex of the form $(1, \dots, 1, \rho_{i-1}(h), \rho_i(h'), \rho_{i+1}(h''), \dots)$ with $h \neq 1$, define the g -image of this vertex as $(1, \dots, 1, \rho_{i-1}(h), \rho_i(gh'), \rho_{i+1}(h''), \dots)$.

Adding to the disjoint union $T_C(G) \cup T_S(G)$ an additional root $\emptyset$ together with edges to the roots of these trees (denoted by $\emptyset_C$ and $\emptyset_S$) we obtain the required tree T . Since the G -set $T_C(G)$ is equivariantly embedded into T the latter is a faithful G -set.

For any subset $J = \{j_1, j_2, \dots\} \subset \{1, 2, \dots\}$ consider the sequence $g_{j_1} \cdot g_{j_2} \cdot \dots \cdot g_{j_k}$, $k \in \omega$. Since G is compact, there is the limit g_J of this sequence. By the definition

of the action of G on $T_S(G)$ we easily see that for $J \neq K$ the elements g_J and g_K have distinct isometry types.

(2) The proof of this statement is similar to the proof above. We modify it as follows. Replace $T_C(G)$ above by any faithful representation of G on the rooted tree $T(p)$ of valency $p + 1$ (see Proposition 2 of [13]). It is well-known that any finite p -group can be embedded into the isometry group of a finite part of $T(p)$ (consisting of several levels). Thus every $\rho_i(G)$ can be presented as an isometry group of some $T(p)_{k_i}$ (consisting of k_i levels). We now replace $T_S(G)$ above by the tree $T(p)$. We define an action of G on this tree as follows. For every $g \in G$ and every vertex of the form $(0^{k_1}, \dots, 0^{k_{i-1}}, 1, t_1, t_2, \dots, t_{k_i}, \dots)$ define the g -image of this vertex as $(0^{k_1}, \dots, 0^{k_{i-1}}, 1, \rho_i(g)t_1, \rho_i(g)t_2, \dots, \rho_i(g)t_{k_i}, \dots)$ (we define the action of g identically on other elements). The rest is obvious. $\square$

Theorem 2.8. *Let G be a nilpotent profinite group. If all representations of G in isometry groups of locally finite trees have small numbers of isometry types, then G is a group of finite rank.*

Proof. Assume that the rank of G is infinite. By Theorem 2.4 (1) there is a prime number p such that the Sylow p -subgroup of G has infinite rank. Assume that G is a pro- p -group. By composing trees as in the proof of Lemma 2.7 we can reduce the situation to this case.

Take the upper central series $1 = G_0 < G_1 < \dots < G_n = G$. It is clear that all G_i are closed. Since $rk(G) = \infty$ we have some i with infinite $rk(G_{i+1}/G_i)$. Find the least $i \leq n$ such that all $rk(G_{i+j+1}/G_{i+j})$ are finite. We denote this number by m . Thus $rk(G/G_m)$ is finite and the group G_m/G_{m-1} is a $\mathbb{Z}_p$ -module which does not have a finite generating set.

Let cosets $g'_1 G_m, \dots, g'_t G_m$ generate topologically G/G_m and K be the intersection of G_m with the closure of the subgroup $\langle g'_1, \dots, g'_t \rangle$. Note that the closure of $\langle g'_1, \dots, g'_t \rangle G_{m-1}$ is a normal subgroup of G such that the corresponding quotient is isomorphic to G_m/KG_{m-1} ($= cl(\langle g'_1, \dots, g'_t \rangle)G_m / cl(\langle g'_1, \dots, g'_t \rangle)G_{m-1}$). Thus to prove the statement it suffices to build a sequence $g_1, g_2, \dots, g_i, \dots \in G \setminus G_{m-1}$ representing pairwise distinct cosets of G_m/KG_{m-1} which satisfy the conditions of Lemma 2.7(1) with respect to this group.

Assume that $g_1, g_2, \dots, g_l$ and the corresponding ρ_i are already defined. Since G_m/KG_{m-1} is a $\mathbb{Z}_p$ -module which does not have a finite generating set, there is an open normal subgroup L of G_m containing KG_{m-1} such that $L/KG_{m-1} < (\bigcap_{i=1}^l \ker(\rho_i)) \cap \langle g_1, \dots, g_l \rangle L$. Let g_{l+1} represent an L -coset of $\bigcap_{i=1}^l \ker(\rho_i)/L$ which does not belong to $\langle g_1, \dots, g_l \rangle L$ and let ρ_{l+1} be the natural homomorphism onto

$$(\bigcap_{i=1}^l \ker(\rho_i)) / (\bigcap_{i=1}^l \ker(\rho_i) \cap \langle g_1, \dots, g_l \rangle L).$$

It is easy to see that this construction produces a required sequence. $\square$

Theorem 2.9. (1) *Let Δ be a set of prime numbers. For each $p_i \in \Delta$ let G_i be a pro- p_i -group such that all faithful representations of G_i in isometry groups of locally finite rooted trees have a small number of isometry types. The set Δ is finite exactly when every faithful representation of $\bigoplus_{p_i \in \Delta} G_i$ in the isometry group of a locally finite rooted tree has a small number of isometry types.*

(2) *An abelian profinite group G has a small number of isometry types for all faithful representations of G in isometry groups of locally finite rooted trees if and only if the set of prime numbers dividing $|G|$ is finite and G is of finite rank.*

Proof. (1) Necessity of the statement follows from Theorem 2.4. Consider the case of a representation of $H \oplus G_s$, where H is a pro- Δ -group where Δ is finite and $p_s \notin \Delta$. Assume that all representations of H as closed subgroups of isometry groups of locally finite rooted trees have a small number of isometry types. We want to prove that so does the group $H \oplus G_s$.

Let $H \oplus G_s$ act on T as a closed subgroup of $\text{Iso}(T)$. Let $g \in G_s$. Consider the tree of g -cycles $T/\langle g \rangle$ induced by g on T . To each vertex $\langle g \rangle t$ of $T/\langle g \rangle$ we assign two numbers: $v(t)$, the valency of t , and $m(t) = |\langle g \rangle t|$. Since G_s has a small number of isometry types, there is a small number of such trees corresponding to elements of G_s (up to label preserving isomorphism). Note that $m(t)$ is a power of p_s .

Since $[H, G_s] = 1$ the action of H on T induces an action of H on $T/\langle g \rangle$. If $h \in H$ defines a h -cycle $\langle h \rangle t$ in T , then let $l_h(t)$ be the length of the corresponding h -cycle in $T/\langle g \rangle$. Since p_s does not divide $|\langle h \rangle t|$, we see that $l_h(t) = |\langle h \rangle t|$. Moreover we also see that the length of the gh -cycle of t is equal to $m(t)l_h(t)$. Now it

is straightforward that the isometry type of h on $T/\langle g \rangle$ determines the isometry type of gh on T (even the isometry type of the pair (g, h)). Since the number of isometry types of elements of H on $T/\langle g \rangle$ is less than continuum we easily obtain the statement of the proposition.

(2) Let Δ be a set of prime numbers. Let $G = \prod_{p \in \Delta} \mathbb{Z}_p^{l_p}$ where all l_p are finite. It is enough to prove that if Δ is finite then every faithful representation of G in $Iso(T)$ has a small number of isometry types.

By part (1) it suffices to consider the case when Δ is a singleton. Let $l_p = l \geq 1$. Let T be a tree as in the formulation and $\mathbb{Z}_p^l$ act as a closed subgroup of $Iso(T)$. By π_n we denote the canonical homomorphism from $\mathbb{Z}_p^l$ to the group of all restrictions of $\mathbb{Z}_p^l$ to T_n , $n \in \omega$. Thus for every $g \in \mathbb{Z}_p^l$ every π_n maps the group $g^{\mathbb{Z}_p}$ to a cyclic p -group.

We fix a tuple of free topological generators of $\mathbb{Z}_p^l$: $g_1, \dots, g_l$ (so that $\mathbb{Z}_p^l = g_1^{\mathbb{Z}_p} \times \dots \times g_l^{\mathbb{Z}_p}$). To an element $g_1^{z_1} \dots g_l^{z_l}$ we associate the tuple $(v_p(z_1), \dots, v_p(z_l))$ of the corresponding values with respect to the p -adic valuation. We claim that given an action of $\mathbb{Z}_p^l$ on T the tuple $(v_p(z_1), \dots, v_p(z_l))$ determines the size of the $g_1^{z_1} \dots g_l^{z_l}$ -cycle of any $t \in T$. In particular this tuple determines the isometry type of $g_1^{z_1} \dots g_l^{z_l}$ (thus the action has a small number of isometry types). We prove this by induction by l .

Let $l = 1$. For any $t \in T_n$ the $g_1^{z_1}$ -cycle of t is determined by $\pi_n(g_1^{z_1})$. Thus to find $|g_1^{z_1}t|$ we may assume that z_1 belongs to $\mathbb{Z}$. Let $v_p(z_1) = s$, i.e. $z_1 = p^s k$ with $(k, p) = 1$. Since the g_1^k -cycle of t coincides with the g_1 -cycle of t , the length of the $g_1^{z_1}$ -cycle of t equals the first integer not less than $|\langle g_1 \rangle t|/p^s$. The rest of this case is an obvious application of [10].

Now consider the case of an action of $H \times g_l^{\mathbb{Z}_p}$, where $H = g_1^{\mathbb{Z}_p} \times \dots \times g_{l-1}^{\mathbb{Z}_p}$. Assume that for any isometric action of H on a locally finite rooted tree T' and any $t' \in T'$ the number $|\langle h \rangle t'|$, $h \in H$, is determined by the corresponding tuple of p -adic values of h . In particular the isometry type of h is determined by this tuple of p -adic values. We want to prove these statement for the action on T and any $g_1^{z_1} \dots g_l^{z_l}$.

As above we assume that $t \in T_n$ and $z_l = p^s k$ with $(k, p) = 1$. Then g_l and g_l^k have the same cycles of t . Let $h = g_1^{z_1} \dots g_{l-1}^{z_{l-1}}$. Consider the tree of h -cycles

$T/\langle h \rangle$ induced by h on T . To each vertex $\langle h \rangle t'$ of $T/\langle h \rangle$ we assign two numbers: $v(t')$, the valency of t' , and $m(t') = |\langle h \rangle t'|$. Note that $m(t')$ is a power of p and is determined by the corresponding tuple of p -adic values of h .

Since $[H, g_l] = 1$ the action of g_l on T induces an action of g_l on $T/\langle h \rangle$. Then g_l and g_l^k have the same cycles of $\langle h \rangle t$.

As a result we see that $|\langle g_l^{z_l} \rangle t|$ equals the first integer not less than $|\langle g_l \rangle t|/p^s$, and the length of the $g_l^{z_l}$ -cycle of $\langle h \rangle t$ in $T/\langle h \rangle$ equals the first integer not less than $|\langle g_l \rangle (\langle h \rangle t)|/p^s$. Dividing $|\langle g_l^{z_l} \rangle t|$ by the length of the $g_l^{z_l}$ -cycle of $\langle h \rangle t$ we find $|(\langle h \rangle t) \cap (\langle g_l^{z_l} \rangle t)|$. This number uniquely determines all equalities $h^i(t) = g_l^{j z_l}(t)$ for $i \leq |\langle h \rangle t|$, $j \leq |\langle g_l^{z_l} \rangle t|$. In particular it uniquely determines the length of the $h g_l^{z_l}$ -cycle of t . $\square$

Conjecture. Let G be a closed subgroup of $GL_d(\mathbb{Z}_p)$. Then G has a small number of isometry types for all faithful actions on locally finite rooted trees.

In the case of abelian groups the conjecture holds by Theorem 2.9 (2). By Lemma 2.7 we see that groups of the form H^ω always have a representation with continuum of isometry types. On the other hand it is easy to see that this lemma cannot be applied to closed subgroups of $GL_d(\mathbb{Z}_p)$. Thus our conjecture looks reasonable. Another interesting question is to prove the converse of Theorem 2.8.

3. ROOTED GROUPS INTERPRETABLE IN $\mathbb{Q}_p$

One of the main obstacles to deeper study of isometry groups of rooted trees is the fact that there is no interesting first-order structure where these groups can be interpreted. To remedy this situation one can choose a rich algebraic object and consider isometry groups interpretable in it. Then the general case can be somehow approximated by information obtained in this way. We now describe one of possible realizations of this idea.

Let T be a locally finite rooted tree. Let Γ be a closed subgroup of $Iso(T)$. We say that $g, g' \in Iso(T)$ have the same Γ -type if there exists $h \in \Gamma$ such that $g^h = g'$. It makes sense to study Γ -types of $Iso(T)$ assuming that Γ is interpretable in some natural model-theoretic objects. In this situation we say that $G \leq Iso(T)$ has a

small number of Γ -types if G meets $\leq \aleph_0$ Γ -orbits of the conjugacy action of Γ on $Iso(T)$.

In this section we concentrate on trees T and groups $\Gamma \leq Iso(T)$ which are interpretable in the theory of the p -adics $\mathbb{Q}_p$. This provides new tools and, moreover, gives a possibility of extension of the matter of the previous sections by some additional questions.

For every natural n we introduce the following equivalence relation on $Iso(T)$: for $g, g' \in Iso(T)$ we define $g \equiv_n^\Gamma g'$ if the restrictions of g and g' to T_n are conjugated by an element of the projection of Γ to T_n . For a subgroup $G \leq Iso(T)$ we denote by c_n^G the number of all $\equiv_n^\Gamma$ -classes meeting G . We will also study the question when the zeta function $\sum_{n \geq 0} c_n^G t^n$ is rational. Applying the main results of [9] we give several positive examples.

We start with the group $GL_d(\mathbb{Z}_p)$ and the corresponding tree T_{ad} , i.e. the tree of cosets in $(\mathbb{Z}_p)^d$ with respect to all subgroups of the form $p^k(\mathbb{Z}_p)^d$.

Let Γ be a closed subgroup of $GL_d(\mathbb{Z}_p)$. Then the group $D = (\mathbb{Z}_p)^d$ becomes a continuous $\Gamma\mathbb{Z}_p$ -module. Thus Γ acts by isometries on T_{ad} . We now give some examples which show that typical subgroups of $GL_d(\mathbb{Z}_p)$ have continuum Γ -types with respect to their standard action on T_{ad} .

Example 1. Let $\Gamma = GL_d(\mathbb{Z}_p) = G$ with $d > 1$. Since $|\mathbb{Z}_p| = 2^{\aleph_0}$, one can produce continuum many Jordan matrices over $\mathbb{Z}_p$.

Assuming that $d = 2$ and G is the group $SD_2(\mathbb{Z}_p)$ of all diagonal matrices over $\mathbb{Z}_p$ with determinant 1 we obtain an abelian group of isometries of the tree T_{ad} having continuum Γ -types where $\Gamma = GL_d(\mathbb{Z}_p)$.

On the other hand extending this idea we can prove the following statement.

Proposition 3.1. *Let $\Gamma = GL_d(\mathbb{Z}_p)$ and $G \leq GL_d(\mathbb{Z}_p)$. If G is a closed subgroup of $Iso(T_{ad})$ which does not have a soluble subgroup of finite index, then G meets $2^{\aleph_0}$ Γ -types.*

Proof. By Tits alternative G has a free subgroup. The proof of [20] shows that G contains a semisimple element g of infinite order. Then any pair of powers $g^{\mathbb{Z}_p}$ cannot be conjugated in Γ . $\square$

Example 2. Let $\Gamma = SL_2(\mathbb{Z}_p)$ and $G = UT_2(\mathbb{Z}_p)$ be the abelian group of all unitriangular matrices over $\mathbb{Z}_p$, $p > 2$. Straightforward computations show that matrices $(a_{11}, a_{12}, a_{21}, a_{22})$ and $(b_{11}, b_{12}, b_{21}, b_{22})$ from G (written as vectors, i.e. $a_{21} = b_{21} = 0$ and $a_{11} = b_{11} = a_{22} = b_{22} = 1$) are conjugated by $(x_{11}, x_{12}, x_{21}, x_{22}) \in \Gamma$ if and only if $x_{21} = 0$ and $b_{12} = x_{11}^2 a_{12}$.

It is worth noting that since $(\mathbb{Z}_p^\times)^2 = \mathbb{Z}_p^\times$, any two elements $\bar{a}, \bar{b} \in G$ as above with $\nu(a_{12}) = \nu(b_{12})$, are Γ -conjugated (by a diagonal matrix). In this case G has a small (i.e. countable) number of Γ -types.

Remark. It is natural to ask if the actions above have a small number of isometry types with respect to the corresponding tree T and $Iso(T)$. It is clear that if for some $\Gamma < Iso(T)$, G has a small number of Γ -types then G has a small number of isometry types. Thus the case when $G = UT_2(\mathbb{Z}_p)$ acts on the corresponding tree T_{ad} is clear.

Assume that G is the group $SD_2(\mathbb{Z}_p)$ of all diagonal matrices over $\mathbb{Z}_p$ with determinant 1. Then

the natural isometric action of G on the tree T_{ad} has a small number of isometry types.

Indeed identify G with the set of diagonals $\{(u, u^{-1}) : u \in \mathbb{Z}_p^\times\}$. Then G acts on the module $\mathbb{Z}_p \times \mathbb{Z}_p$ by the action:

$$(u, u^{-1}) \cdot (z_1, z_2) = (uz_1, u^{-1}z_2).$$

This action naturally induces the action of G on T_{ad} . It is clear that $G \cong \mathbb{Z}_p^\times$. On the other hand it is known that the group $\mathbb{Z}_p^\times$ can be decomposed into a direct sum isomorphic to $\mathbb{Z}(p) \oplus \mathbb{Z}_p$, where the factor isomorphic to $\mathbb{Z}(p)$ is represented by the cyclic subgroup $\{\varepsilon_1, \dots, \varepsilon_{p-1}\}$ of $p-1$ -th roots of unity. The factor $\mathbb{Z}_p$ is represented by all elements of $\mathbb{Z}_p^\times$ which are congruent to 1 mod p (see Chapter 18 in [7]). It only remains to apply Theorem 2.9.

In fact the latter factor contains l such that all other elements are represented by l^z , where $z \in p\mathbb{Z}_p$. To find this l let t be a primitive root mod p which is a primitive root modulo every power of p and let $l = t^{p-1}$ (see [7], p.316).

As a result the action of $(u, u^{-1}) = (l^z, l^{-z})$ on a vertex $(z_1, z_2) + p^k \mathbb{Z}_p$ is defined as follows. Assuming that $z_1, z_2 \in \omega$ and $z_1, z_2 < p^k$ we represent $(z_1, z_2) =$

$(p^{s_1}\varepsilon_i l^{t_1}, p^{s_2}\varepsilon_j l^{t_2})$ with $s_1, s_2 < k$ and $t_1 < k - s_1 - 1$ and $t_2 < k - s_2 - 1$ (this is possible because $l^i \equiv 1 \pmod{p^k}$ if and only if $p^{k-1} \mid i$). Then the action $(u, u^{-1}) \cdot ((z_1, z_2) + p^k \mathbb{Z}_p)$ is defined by the representative $(p^{s_1}\varepsilon_i l^{z+t_1}, p^{s_2}\varepsilon_j l^{-z+t_2})$.

According to Proposition 3.1, when Γ and T_{ad} are as in this proposition, groups with a small number of Γ -types must be almost soluble. According to Theorem 1 of [12] the Hilbert-Poincaré series of such a group $G < GL_d(\mathbb{Z}_p)$ is $\mathbb{Q}$ -rational.

To show that zeta-functions of the numbers of conjugacy classes are rational for these examples we now formulate some general theorem.

We consider the p -adics $\mathbb{Q}_p$ with the standard valuation $\nu : \mathbb{Q}_p^* \rightarrow \mathbb{Z}$ (and the value group), the valuation ring $\mathcal{O} = \mathbb{Z}_p$, the maximal ideal $M = \mathbb{Z}_p \setminus \mathbb{Z}_p^\times$ and the corresponding residue field. We remind the reader that ν is a homomorphism such that $\nu(x + y) \geq \inf(\nu(x), \nu(y))$ and $\nu(p) = 1$. Then $\mathbb{Z}_p = \{x \in \mathbb{Q}_p : \nu(x) \geq 0\}$ is the valuation ring of $\mathbb{Q}_p$ and $\mathbb{Z}/p\mathbb{Z}$ is the corresponding residue field.

For each $n \in \omega$ we add the sort $GL_n(\mathbb{Q}_p)$ and the set of lattices $S_n(\mathbb{Q}_p) = GL_n(\mathbb{Q}_p)/GL_n(\mathbb{Z}_p)$. We also add the natural map $GL_n(\mathbb{Q}_p) \rightarrow S_n(\mathbb{Q}_p)$. Structures of this form were introduced in [8] and [9], where sorts of these structures were called geometric. It is proved in [9] that $\mathbb{Q}_p$ in this language admits elimination of imaginaries. The following theorem from [9] (Theorem 6.2) plays the central role in this section.

Let $R = (R_l)_{l \in \omega^r}$ be a definable family of subsets of $\mathbb{Q}_p^N$ (i.e. a definable subset of $\mathbb{Q}_p^N \times \mathbb{Z}^r$, where $\mathbb{Z}$ is the value group and $\omega \subset \mathbb{Z}$). Let $E = (E_l)_{l \in \omega^r}$ be a definable family of equivalence relations on R (i.e. every equivalence class is contained in an l -fibre of R for some $l \in \omega^r$). Suppose that for each $l \in \omega^r$ the set of equivalence classes R_l/E_l is finite. Let $a_l = |R_l/E_l|$. Then the power series $\sum_{l \in \omega^r} t^l \in \mathbb{Q}[[t_1, \dots, t_r]]$ is $\mathbb{Q}$ -rational.

We now describe how it can be applied in our situation.

Proposition 3.2. *Let Γ and G be subgroups of $GL_d(\mathbb{Z}_p)$ definable in $\mathbb{Q}_p$. Let c_n^G be the number of all $\equiv_n^\Gamma$ -classes meeting G defined with respect to T_{ad} . Then the zeta function $\sum_{n \geq 0} c_n^G t^n$ is $\mathbb{Q}$ -rational.*

Proof. We interpret T_{ad} in $\mathbb{Q}_p$ as the set of equivalence classes on $\mathbb{Z}_p^d \times \omega$ (where $\omega \subset \mathbb{Z}$ is the non-negative part of the value group) with respect to the following equivalence relation. Let

$$(\bar{z}_1, c_1) \sim (\bar{z}_2, c_2) \text{ if } (c_1 = c_2) \wedge (\nu(\bar{z}_1 - \bar{z}_2) \geq c_1).$$

It is clear that this defines the set of cosets of all groups of the form $p^c(\mathbb{Z}_p)^d$. The inclusion relation of cosets (which is the order relation of T_{ad}) is defined by representatives:

$$(\bar{z}_1, c_1) \leq (\bar{z}_2, c_2) \Leftrightarrow (c_1 \geq c_2) \wedge (\nu(\bar{z}_1 - \bar{z}_2) \geq c_1).$$

Let $R = (R_i)_{i \in \omega} = G \times \omega$. We define a family $E = (E_i)$ of equivalence relations on R as follows:

$$((g_1, c_1), (g_2, c_2)) \in E \Leftrightarrow (c_1 = c_2) \wedge (\exists \gamma \in \Gamma)$$

$$(\forall \bar{z} \in (\mathbb{Z}_p)^d)(\forall c \leq c_1)(\gamma(g_1([\bar{z}, c])_\sim) = g_2(\gamma([\bar{z}, c])_\sim)).$$

The rest follows by Theorem 6.2 of [9]. $\square$

Proposition 3.3. *Let G be a finitely generated, torsion free, nilpotent pro- p -group. Then for some natural d there is an embedding of G into $GL_d(\mathbb{Z}_p)$ such that the corresponding action (G, T_{ad}) is definable in $\mathbb{Q}_p$. The zeta function $\Sigma_{n \geq 0} c_n^G t^n$ of this action with respect to $\Gamma = GL_d(\mathbb{Z}_p)$ is $\mathbb{Q}$ -rational.*

Proof. Let $a_1, \dots, a_n$ be a Malcev basis of G (i.e. any element of G can be written uniquely in the form $a_1^{\lambda_1} \cdot \dots \cdot a_n^{\lambda_n}$, $\lambda_i \in \mathbb{Z}_p$). A well-known theorem (see [17]) states that group multiplication and inversion in G are given by polynomials in λ_i with coefficients in $\mathbb{Q}$ as is the map $G \times \mathbb{Z}_p \rightarrow G$, $(g, \lambda) \rightarrow g^\lambda$. This gives an interpretation of G in $\mathbb{Q}_p$ as $\mathbb{Z}_p^n$.

By Theorem 59.2.1 from [17] there is a natural number d and a polynomial map $\phi : \mathbb{Z}_p^n \rightarrow UT_d(\mathbb{Z}_p)$ giving an isomorphic representation of G on $\mathbb{Q}_p$. Moreover the converse map ϕ^{-1} is linear. This clearly provides a definable action (G, T_{ad}) . By Proposition 3.2 we have the last statement of our proposition. $\square$

One of the main examples of G -trees in geometric group theory is the construction of $SL_2(\mathbb{Q}_p)$ -tree of lattices of Serre and Tits. We now show that the actions

studied in the proposition above (and in Proposition 3.2) arise as point-stabilizers in this example. The following well-known construction is taken from [18]).

A *lattice* in $\mathbb{Q}_p \times \mathbb{Q}_p$ is a two-generated $\mathbb{Z}_p$ -submodule of $\mathbb{Q}_p \times \mathbb{Q}_p$ which generates the $\mathbb{Q}_p$ -vector space $\mathbb{Q}_p \times \mathbb{Q}_p$. By $\mathcal{X}$ we denote the set of equivalence classes of lattices with respect to the equivalence

$$L \sim L' \leftrightarrow (\exists z \in \mathbb{Q}_p^*)(L' = Lz).$$

The distance $d([L]^\sim, [L']^\sim)$ between the corresponding classes is defined as follows. Find $L'' \sim L'$ such that $L'' = \langle e_1 p^a, e_2 p^b \rangle < \langle e_1, e_2 \rangle = L$. Then the number $|a - b|$ does not depend on the choice of the basis e_1, e_2 and the representative L'' ; so it is taken as the corresponding distance.

The metric space $(\mathcal{X}, d)$ is a simplicial tree of valency $p + 1$, and $PGL_2(\mathbb{Q}_p)$ has an isometric action on $\mathcal{X}$. Moreover the group $PSL_2(\mathbb{Q}_p)$ acts on $\mathcal{X}$ without inversions. To interpret this G -space in $Th(\mathbb{Q}_p)$, the set of lattices is usually identified with the left coset space $S_2 = GL_2(\mathbb{Q}_p)/GL_2(\mathbb{Z}_p)$ (as above, see [8]). Then $\mathcal{X}$ can be considered as the coset space $GL_2(\mathbb{Q}_p)/(\mathbb{Q}_p^* \cdot GL_2(\mathbb{Z}_p))$, where $PGL_2(\mathbb{Q}_p)$ ($PSL_2(\mathbb{Q}_p)$ resp.) acts from the left and $\mathbb{Q}_p^*$ now stands for the subgroup of scalar matrices.

Formally we consider $\mathcal{X}$ as $GL_2(\mathbb{Q}_p)$ with respect to the following equivalence relation:

$$\begin{aligned} (a_{ij})_{i \leq 2, j \leq 2} \sim (b_{ij})_{i \leq 2, j \leq 2} &\Leftrightarrow (\exists (c_{ij})_{i \leq 2, j \leq 2}, (d_{ij})_{i \leq 2, j \leq 2} \in GL_2(\mathbb{Q}_p)) (\exists q \in \mathbb{Q}_p^*) \\ &[\nu(c_{ij}) \geq 0 \wedge \nu(d_{ij}) \geq 0 \wedge ((c_{ij})_{i \leq 2, j \leq 2} \cdot (d_{ij})_{i \leq 2, j \leq 2} = E) \wedge \\ &((a_{ij})_{i \leq 2, j \leq 2} = (b_{ij})_{i \leq 2, j \leq 2} \cdot (q \cdot c_{ij})_{i \leq 2, j \leq 2})]. \end{aligned}$$

The group $PGL_2(\mathbb{Q}_p)$ ($PSL_2(\mathbb{Q}_p)$ resp.) is considered as $GL_2(\mathbb{Q}_p)$ ($SL_2(\mathbb{Q}_p)$) with respect to the equivalence relation induced by the center. The action of $PGL_2(\mathbb{Q}_p)$ on $\mathcal{X}$ is defined by left multiplication of the corresponding representatives.

To define the distance d on $\mathcal{X}$ we use the following fomula:

$$\begin{aligned} d((a_{ij})_{i \leq 2, j \leq 2}, (b_{ij})_{i \leq 2, j \leq 2}) = k &\Leftrightarrow (\exists t \in \mathbb{Q}_p^*)(\exists (c_{ij})_{i \leq 2, j \leq 2} \in GL_2(\mathbb{Z}_p)) \\ (\exists q, r \in \mathbb{Z}_p)[k = |\nu(q) - \nu(r)| \wedge (qa_{11}, ra_{12}, qa_{21}, ra_{22}) &= (b_{ij})_{i \leq 2, j \leq 2} \cdot (t \cdot c_{ij})_{i \leq 2, j \leq 2}]. \end{aligned}$$

Let $L_0 = \langle (1, 0), (0, 1) \rangle$. The stabilizer of the point $[L_0]^\sim$ is exactly the subgroup of $PGL_2(\mathbb{Q}_p)$ consisting of all cosets of the form $g\mathbb{Q}_p^*$ where $g \in GL_2(\mathbb{Z}_p)$.

Each vertex of $\mathcal{X}$ is represented by a unique lattice $L \subseteq L_0$ such that $L_0/L \cong \mathbb{Z}_p/p^n\mathbb{Z}_p$, where n is the distance between L_0 and L . In this case L/p^nL_0 is a direct factor of rank 1 of the $\mathbb{Z}_p/p^n\mathbb{Z}_p$ -module L_0/p^nL_0 . When $L = \langle u, vp^n \rangle$ for some u and v generating L_0 , we identify L with the $\mathbb{Z}_p/p^n\mathbb{Z}_p$ -module generated by $u + p^nL_0$.

Formally we present the set of these modules as follows. In the rooted tree $\mathcal{L}$ of all cosets of $L_0 = \mathbb{Z}_p \times \mathbb{Z}_p$ of the form $u + p^n(\mathbb{Z}_p \times \mathbb{Z}_p)$ with $\nu(u) = 0$ we introduce the following equivalence relation :

$$u + p^nL_0 \approx u' + p^lL_0 \Leftrightarrow (n = l) \wedge (\exists x \in \mathbb{Z}_p^\times)(xu - u' \in p^nL_0).$$

It is easy to see that the action of any $g\mathbb{Q}_p^*$ with $g \in GL_2(\mathbb{Z}_p)$ on the set of elements of $\mathcal{X}$ which are n -distant from L_0 , corresponds to the action of g on the set of these classes. Moreover if g and $g' \in GL_2(\mathbb{Z}_p)$ determine the same element of $PGL_2(\mathbb{Z}_p)$, then $g(u + p^nL_0) \approx g'(u + p^nL_0)$, where u as above. Thus the stabilizer of the element L_0 in $\mathcal{X}$ is identified with $PGL_2(\mathbb{Z}_p)$, which is considered with respect to its action on $\mathcal{L}/\approx$.

Note that the rooted tree $\mathcal{L}/\approx$ is divided into levels represented by cosets $u + p^nL_0$ with the same p^n . Moreover the vertices $u + p^nL_0$ and $u' + p^{n+1}L_0$ are adjacent if and only if $(\mathbb{Z}_p/p^n\mathbb{Z}_p)(u + p^nL_0) = (\mathbb{Z}_p/p^n\mathbb{Z}_p)(u' + p^nL_0)$ (by the definition of $\mathcal{X}$).

Theorem 3.4. *In the notation above let $\Gamma = G = PGL_2(\mathbb{Z}_p)$.*

(1) *Let c_n^G be the number of all $\equiv_n^\Gamma$ -classes meeting G defined with respect to the rooted tree $\mathcal{L}/\approx$. Then the zeta function $\sum_{n \geq 0} c_n^G t^n$ is $\mathbb{Q}$ -rational.*

(2) *The group G has a small number of isometry types with respect to the action on $\mathcal{L}/\approx$ and there is continuum Γ -types meeting G .*

Proof. (1) Repeating the proof of Proposition 3.2 we interpret $\mathcal{L}/\approx$ in $\mathbb{Q}_p$ as the set of equivalence classes on a definable subset of $\mathbb{Z}_p^2 \times \omega$ (where $\omega \subset \mathbb{Z}$ is the non-negative part of the value group). Then we consider $R = (R_i)_{i \in \omega} = GL_2(\mathbb{Z}_p) \times \omega$, with respect to the family $E = (E_i)$ of equivalence relations on R as it was described above. The rest follows by Theorem 6.2 of [9].

(2) By obvious arguments from linear algebra we may assume that elements of $PGL_2(\mathbb{Z}_p)$ are presented by upper triangular matrices. Let (a_{11}, a_{12}, a_{22}) be

a tuple of non-zero entries of such a matrix $g \in G$. Then multiplying g^n with $u = (z_1, z_2) \in \mathbb{Z}_p \times \mathbb{Z}_p$ we obtain a representative of the projective line corresponding to

$$(z_1 + \frac{a_{12}z_2}{a_{11} - a_{22}}(1 - (\frac{a_{22}}{a_{11}})^n), (\frac{a_{22}}{a_{11}})^n z_2).$$

Decompose the group $\mathbb{Z}_p^\times$ into a direct sum isomorphic to $\mathbb{Z}(p) \oplus \mathbb{Z}_p$, where the factor isomorphic to $\mathbb{Z}(p)$ is represented by the cyclic subgroup $\{\varepsilon_1, \dots, \varepsilon_{p-1}\}$ of $p-1$ -th roots of unity. The factor $\mathbb{Z}_p$ is represented by all elements of $\mathbb{Z}_p^\times$ which are congruent to 1 mod p (see Chapter 18 in [7]). As above we find l such that all elements of this factor are represented by l^z , where $z \in p\mathbb{Z}_p$ (see [7], p.316). Assuming that $a_{22}/a_{11} = \varepsilon_j l^z$ we see that $(\frac{a_{22}}{a_{11}})^n \equiv 1 \pmod{p^k}$ if and only if $\varepsilon_j^n = 1$ and $p^{k-1} | nz$.

Assume $(a'_{11}, a'_{12}, a'_{22})$ be a tuple of non-zero entries of a matrix $g' \in G$ such that $\nu(a_{12}) = \nu(a'_{12})$, $a'_{22}/a'_{11} = \varepsilon_j l^{z'}$ and $\nu(z) = \nu(z')$. Then for every pair $(z_1, z_2) \in \mathbb{Z}_p \times \mathbb{Z}_p$ the elements g and g' have cycles of the same length at the element of $\mathcal{X}$ represented by $(z_1, z_2) + p^m L_0$. This proves the statement. $\square$

Remark. There is another version of Proposition 3.2 which does not use definability in $\mathbb{Q}_p$ and which can be also applied to examples of this kind.

Proposition 3.5. *Let Γ and G be closed subgroups of $GL_d(\mathbb{Z}_p)$. Let c_n^G be the number of all $\equiv_n^\Gamma$ -classes meeting G defined with respect to T_{ad} . Then the zeta function $\sum_{n \geq 0} c_n^G t^n$ is $\mathbb{Q}$ -rational.*

To prove this proposition we must apply a more involved argument. In fact we must repeat the proof of Theorem 1.2 from [4] with some slight changes. Since this theorem is outside the main theme of the paper we just mention that our adaptation requires that formula (1.1) from [4] (counting the number of conjugacy classes in a finite group) must be replaced by the formula

$$c_n^G = |\Gamma_n|^{-1} \sum \{ |C_{\Gamma_n}(g)| : g \in GL_d(\mathbb{Z}_p/p^n \mathbb{Z}_p) \\ \wedge \exists g' \in G_n (g \equiv_n^\Gamma g') \},$$

where G_n and Γ_n are the natural projections of G and Γ with respect to $GL_d(\mathbb{Z}_p/p^n \mathbb{Z}_p)$. Since any closed subgroup of $GL_d(\mathbb{Z}_p)$ is compact p -adic analytic, the arguments

of Section 1 of [4] show that G , Γ and $\equiv_n^\Gamma$ are definable in the theory of analytic functions from [2]. This allows us to adapt the proof from [4] to our case.

4. BRANCH GROUPS. MODEL THEORETIC ASPECTS.

As we have noted above, branch groups have a large number of isometry types. Can they be defined (in $\mathbb{Q}_p$) as closed subgroups of matrices over $\mathbb{Z}_p$? This is not true. Indeed, let G be the profinite completion of Grigorchuk's 3-generated 2-group from [11]. In fact it is shown in [12] that the coefficients a_n of the Hilber-Poincaré series of G do not have a polynomial growth. By Theorem 1 of [12] (or Interlude A of [3]) the group G is not a closed subgroups of matrices over $\mathbb{Z}_p$.

How do branch groups look with respect to other model theoretic properties of profinite groups studied so far? In this section we concentrate on ω -categoricity of branch groups in the language introduced by Z.Chatzidakis in [5].

Let G be a profinite group. As in [5] to G we associate a structure $S(G)$ of the language $L = (\leq, \sim, C^2, P^3, 1)$ as follows. The structure $S(G)$ is defined on the set of all cosets gN for all open normal subgroups $N < G$. The symbol P is interpreted as follows:

$$S(G) \models P(g_1N_1, g_2N_2, g_3N_3) \Leftrightarrow (N_1 = N_2 = N_3) \wedge (g_1g_2N_1 = g_3N_1).$$

The symbol C corresponds to inclusion:

$$S(G) \models C(gN, hM) \Leftrightarrow (N \subseteq M) \wedge (gM = hM).$$

The relation $gN \leq hM$ means $N \subseteq M$ and we define $gN \sim hM \Leftrightarrow (gN \leq hM) \wedge (hM \leq gN)$. The constant 1 corresponds to G .

It is convenient to view $S(G)$ as an ω -sorted structure with respect to sorts $S_n = \{gN : |G : N| = n\}$, $n \in \omega$. In this language the class of structures $S(G)$ becomes elementary. In Section 1.5 of [5] some natural axioms of this class are given: it is shown that for any structure S satisfying these axioms the $\sim$ -classes naturally form a projective system of finite groups such that S is $S(G)$ for the projective limit G of this system.

We say that $A \subseteq S(G)$ forms a substructure if $(\forall x, y \in S(G))((x \leq y) \wedge (x \in A) \rightarrow y \in A)$ and $(\forall x, y \in A)(\exists z \in A)((z \leq y) \wedge (z \leq x))$. It is straightforward

that any substructure $A \subseteq S(G)$ defines a profinite group H with $A = S(H)$ such that H is a continuous homomorphic image of G . On the other hand if $\phi : G \rightarrow H$ is a continuous epimorphism of profinite groups, then the map $S(H) \rightarrow S(G) : gN \rightarrow \phi^{-1}(gN)$ defines an embedding of $S(H)$ into $S(G)$ as a substructure. The following property can be taken as the definition of ω -categoricity (see Section 1.6 of [5]).

Definition 4.1. *We say that a countable structure $S(G)$ is ω -categorical if for all $n, j_1, \dots, j_k \in \omega$ the group $\text{Aut}(S(G))$ has finitely many orbits on $(S_{j_1} \cup \dots \cup S_{j_k})^n$ (i.e. $\text{Aut}(S(G))$ is oligomorphic).*

It is easy to see that

for every finitely generated profinite group G each sort $S_i(G)$ is finite (see [3], Proposition 1.6); i.e. $S(G)$ is ω -categorical.

Z.Chatzidakis has noticed in [5] (Theorem 2.3) that when G has the *Iwasawa Property* (IP), the structure $S(G)$ is ω -categorical. We remind the reader that G has (IP) if for every epimorphism $\theta : H \rightarrow K$ of finite groups with $H \in \text{Im}(G)$, and for every epimorphism $\phi : G \rightarrow K$ there is an epimorphism $\psi : G \rightarrow H$ such that $\phi = \theta \cdot \psi$. Note that (IP) is very close to projectivity (which in the case of pro- p -groups (IP) is equivalent to p -freeness, see [6], Chapter 20).

It is worth noting that all closed subgroups of $GL_n(\mathbb{Z}_p)$ have ω -categorical structures $S(G)$. This follows from the fact that they have a finite number of open subgroups of index k , $k \in \omega$ (moreover the corresponding function counting the number of subgroups is polynomial [3]). Thus groups considered in the previous sections usually had ω -categorical structures $S(G)$ (excluding $\mathbb{Z}(p)^\omega$).

In the following proposition we consider *just infinite profinite groups*. This means that each proper continuous homomorphic image of the group is finite. Just infinite branch groups are described in Theorem 4 of [13]. It states that a profinite group with a branch structure $\{L_i, H_i : i \in \omega\}$ is just infinite if and only if the commutator subgroup $[L_i, L_i]$ is of finite index in L_i for all i .

Proposition 4.2. *Let $G < \text{Iso}(T)$ be a weakly branch profinite group such that the restricted stabilizer $Rs(n)$ is level-transitive on every subtree of level number n .*

(1) *Then $S(G)$ is ω -categorical if and only if every sort S_k is finite.*

(2) If G is additionally a branch group which is just infinite, then $S(G)$ is ω -categorical.

Proof. (1) The first statement of the proposition is a straightforward consequence of Theorem 7.5 and Proposition 8.1 of [15]. Indeed each automorphism ϕ of $S(G)$ naturally defines a continuous automorphism of the group G . By [15] under our assumptions, each automorphism of G is induced by conjugation of an isometry h of T such that $G^h = G$. This means that if an open subgroup $N < G$ is the pointwise stabilizer in G of the first l levels of T , then $\phi(N) = N$. Since each open subgroup of G contains such a subgroup N we easily see that all $\text{Aut}(S(G))$ -orbits on $S(G)$ are finite.

(2) Let us show that any open normal subgroup $N < G$ has a finite orbit with respect to automorphisms of G induced by conjugation by isometries of T . Let groups L_i and H_i be defined as in the definition of branch groups (see Section 1.1). Since G is just infinite, by Theorem 4 of [13] the commutator subgroup $[L_i, L_i]$ is of finite index in L_i , and thus $[H_i, H_i]$ is of finite index in G . On the other hand for every n there is a number $l = l(n)$ such that every subgroup of G of index n contains an element which does not fix the subtree T_l pointwise.

Assume $|G : N| = n$ and N is normal in G . Find an element $g \in N$ which does not fix the subtree T_l pointwise for $l = l(n)$. By the proof of Theorem 4 the group N contains the commutator subgroup $[H_{i+1}, H_{i+1}]$. Since the index $|G : [H_{i+1}, H_{i+1}]|$ is finite the number of normal subgroups of G of index n is finite. $\square$

Example. Let G be the profinite completion of Grigorchuk's 3-generated 2-group from [11]. Since G is a finitely generated profinite group, $S(G)$ is ω -categorical. It is worth noting that it is proved in [13] that G is a just infinite branch group.

In the remained part of the section we study ω -categoricity of *wreath branch groups* (see [15]). Since it can happen that such a group is not finitely generated and not just infinite, Proposition 4.2 does not work very well here.

Let $A_i, i \in \omega$, be the sequence of alphabets defining T as in Introduction. Assume that for every n we have chosen a permutation group $P_i \leq \text{Sym}(A_i)$. Then the subgroup of all labellings $\{\gamma(v)\}_{v \in T}$ with $\gamma(v) \in P_{i+1}$ for $v \in \text{layer}_i$ is the *iterated wreath product* of (P_i, A_i) , $i \in \{1, 2, \dots\}$ and is denoted by $\wr_\omega(P_i, A_i)$. It is noticed

in [15] (Theorem 8.2) that this group is a branch group. Thus it is called in [15] a wreath branch group.

Proposition 4.3. (1) *Assume that all groups (P_i, A_i) are simple transitive permutation groups. Then the group $G = \wr_\omega(P_i, A_i)$ has the Iwasawa property (IP) and $S(G)$ is ω -categorical.*

(2) *The group $\text{Iso}(T)$ (i.e. $\wr_\omega(\text{Sym}(A_i), A_i)$) does not have the Iwasawa property (IP) and $S(G)$ is not ω -categorical.*

Proof. (1) This statement follows from the observation that every closed normal subgroup of G is of the form $\text{Ker}\pi_n$ for appropriate n . We think that this fact is folklor, but we give some sketch of it. Let K be a closed normal subgroup of G . Let n is the least natural number such that there is $g \in K \setminus \{1\}$ of depth n , i.e. n is the minimal number such that in the corresponding labelling $\{\gamma(v)\}_{v \in T}$ there is $v \in T_n$ with $\gamma(v) \neq 1$. Then we claim that $K = \text{Ker}\pi_n$.

To see this it is enough to show that for every $l \geq n$ every isometry of the form $\{\delta(v)\}_{v \in T}$ such that T_l has a unique non-trivial $\delta(w)$ and $w \in \text{layer}_l$, is contained in K . Let g and n be as above and $v_0 \in T_n$ witness that g is of depth n . We start with the observation that for every $l \geq n$ there is $h \in K$ of depth l . This h can be chosen as an appropriate commutator $[g, f]$ where $f \in G$ is of depth l . For example, if for all $w \in T_l$ which are below v we have $\gamma(w) = 1$, then we just arrange that the labelling of f is non-trivial only at two points $w, w^g \in T_l$ below v and the corresponding $(\sigma(w))^{-1}$ is not equal to $\sigma(w^g)$.

To finish the proof of our claim take an isometry $g_\delta \in K$ defined by $\{\delta(v)\}_{v \in T}$ as in the previous paragraph. Choose $w \in \text{layer}_l$ with non-trivial $\delta(w)$. Conjugating g_δ by elements representing labellings with the unique non-trivial element at w we generate a subgroup of K having an element g' such that the corresponding labelling is non-trivial at w but trivial at all points of $T_l \setminus \{w\}$. By simplicity of (P_{l+1}, A_{l+1}) , the point w in our g' can be labelled by any element of P_{l+1} . It is also clear that conjugating g' by elements of G we can move w to any element of layer_l . This finishes the proof.

(2) Here we use implicitly the characterization of closed normal subgroups of the group $\text{Iso}(T) = \wr_\omega(\text{Sym}(A_i), A_i)$ from [19]. For every natural l consider the subgroup (denoted by N_l) of all labellings $\{\gamma(v)\}_{v \in T}$ such that the product

$\prod \{sgn(\gamma(v)) : v \in layer_l\}$ is 0. By $sgn(\sigma)$ we denote the parity of the permutation σ (i.e. it equals to 1 when it is odd). It is easy to see that N_l is a normal closed subgroup of $Iso(T)$, $l \in \omega$, and $|Iso(T) : N_l| = 2$. Thus the sort of $S(Iso(T))$ consisting of normal open subgroups of index 2, is infinite. Since $Iso(T)$ is a branch group by Proposition 4.2 we see that $S(Iso(T))$ is not ω -categorical. Thus it does not satisfy (IP). $\square$

Note that the Iwasawa property is slightly casual in the context of branch groups.

Proposition 4.4. *Let $T = T_{\bar{m}}$, where 8 divides some $m_1 \cdot \dots \cdot m_i$. Let $G < Iso(T)$ be a profinite branch group with the corresponding system $\{H_i, L_i : i \in \omega\}$. Then there is a profinite branch group G^* which does not satisfy (IP), such that for some natural numbers k and k' the branch system of G^* with indexes above k becomes $\{H_i, L_i : i \geq k'\}$. Moreover $S(G^*)$ is ω -categorical if and only if $S(G)$ is ω -categorical.*

Proof. To prove this find k , such that the number $l = |layer_k|$ is of the form $8l'$. Consider the tree $T' = T_{\bar{m}'}$ with $m'_1 = l$ and $m'_i = |layer_{k+i-1}|$ for $i \geq 2$. The group G^* acts on T' as follows. Let the kernel $Ker \pi_1(G^*)$ be H_k with respect to the action copied from the action of H_k on the k -th layer of T . Define $\pi_1(G^*)$ to be a regular action of $\mathbb{Z}(2) \oplus \mathbb{Z}(4l')$ on the first layer of T' . Then let G^* act on T' as the wreath product of these groups.

To see that G^* does not satisfy (IP) consider the resulting homomorphism ϕ from

$$G^* \rightarrow \pi_1(G^*) \rightarrow \mathbb{Z}(2),$$

where the second one is the projection to $\mathbb{Z}(2)$. As an epimorphism θ of finite groups $H \rightarrow K$ take $\mathbb{Z}(4l') \rightarrow \mathbb{Z}(2)$ induced by $2l'$ -th exponentiation. If $\theta : G^* \rightarrow \mathbb{Z}(4l')$ makes this diagram commutative, then the generator of $\mathbb{Z}(2)$ in the decomposition $\mathbb{Z}(2) \oplus \mathbb{Z}(4l')$ must have a $2l'$ -root. This is a contradiction.

The second statement of the proposition follows from the fact that for any profinite group H and an open subgroup P the structure $S(H)$ is ω -categorical if and only if $S(P)$ is ω -categorical. This follows from the fact that the map $K \rightarrow P \cap K$ for open subgroups $K < H$ is finite-to-one. $\square$

As we already noted a finitely generated profinite group G has finitely many open subgroups of index n . Thus the structure $S(G)$ for any closed subgroup $G < GL_d(\mathbb{Z}_p)$ or $PGL_d(\mathbb{Z}_p)$, realizes the simplest case. This shows that if any of the groups which we mentioned in the introduction, has ω -categorical $S(G)$, then it becomes very simple. When it is not ω -categorical, then it becomes extremely complicated branch group (very close to the whole $Iso(T)$). This shows that the properties studied in Sections 2 and 3 correspond to our example much better than ω -categoricity of $S(G)$.

REFERENCES

- [1] H.Bass, M.V.Otero-Espinar, D.N.Rockmore and C.P.L.Tresser, Cyclic Renormalization and Automorphism Groups of Rooted Trees. Springer Verlag, 1996.
- [2] J.Denef and L.van den Dries, p -adic and real subanalytic sets, Ann. Math., 128(1988), 79 - 138.
- [3] J.P.Dixon, M.P.F.du Sautoy, A.Mann and D.Segal, Analytic pro- p -groups, 2nd edition, Cambridge Univ. Press, Cambridge, 1999.
- [4] M.P.F.du Sautoy, Counting conjugacy classes, Bull.London Math.Soc. 37(2005), 37 - 44.
- [5] Z.Chatzidakis, Model theory of profinite groups having the Iwasawa property, Illinois Journal of Math., 42(1998), 70 - 96.
- [6] M.D.Fried and M.Jarden, Field Arithmetic, Springer-Verlag, Berlin-Heidelberg, 1986.
- [7] L.Fuchs, Infinite Abelian Groups, I and II, Academic Press, New York - London, 1970, 1973.
- [8] D.Haskell, E.Hrushovski and D.Macpherson, Definable sets in algebraically closed valued fields: elimination of imaginaries, J. Reine und Angew. Math. 597(2006), 175 - 236.
- [9] E.Hrushovski and B.Martin, Zeta functions from definable equivalence relations, ArXiv: math.LO/0701011
- [10] P.Gawron, V.Nekrashevych and V.Sushchansky, Conjugation in tree automorphism groups, Int.J.Alg.Comp.,11(2001), 529 - 547.
- [11] R.Grigorchuk, On the Burnside problem for periodic groups, Func. Anal. Appl. 14(1980), 41 - 43.
- [12] R.Grigorchuk, On the Hilbert-Poincaré series of graded algebras associated with groups, Math. Sb. 180(1989), 207 - 224.
- [13] R.Grigorchuk, Just infinite branch groups. In: New horizons in pro- p -groups (M.P.F.Du Sautoy, D.Segal, A.Shalev, eds), Birkhäuser, Boston, 2000
- [14] K.Krupiński, Products of finite abelian groups as profinite groups, J.Algebra 288(2005), 556 - 582.
- [15] Ya.Lavreniuk and V.Nekrashevych, Rigidity of branch groups acting on rooted trees, Geometria Dedicata 89(2002), 159 - 179.

- [16] L.Newelski, Small profinite groups, J.Symb.Logic, 66(2001), 859 - 872.
- [17] Yu.I.Merzljakov, Rational Groups, Nauka, Moscow, 1987.
- [18] J.-P. Serre, Trees. Springer-Verlag, Berlin-New York, 1980.
- [19] V.Sushchansky, The normal structure of isometry groups of semifinite Baire metrics, In: Infinite Groups and Related Algebraic Structures, 269 - 289, Akad. Nauk Ukrainy, Inst. Mat., Kiev, 1993.
- [20] J. Tits, Free subgroups in linear groups, J. Algebra, 20(1972), 250 - 270.
- [21] F.Wagner, Small profinite m -stable groups, Fund. Math. 176(2003), 181 - 191.
- [22] E.Zelmanov, On periodic compact groups, Israel J.Math. 77(1992), 83 - 95

Institute of Mathematics, Wrocław University,
pl.Grunwaldzki 2/4, 50-384 Wrocław, Poland
e-mail: ivanov@math.uni.wroc.pl